




































Research Samples Visits 2, 6 & unscheduled (baseline, month 12 & as required)

Tubes	Invert tube	Allow to stand	Centrifuge	Transfer	Store	Freeze at -80C
 <p>4-5 mls blood required.</p>	 <p>5 times</p>	 <p>Min 30 minutes Max 60 minutes</p>	 <p>1000g for 15 mins</p>	 <p>MAX 0.5ml per Fludix tube</p>	 <p>Up to 5 Fludix tubes</p>	
 <p>4 mls blood required</p>	 <p>8-10 times</p>	 <p>Max 60 minutes</p>			 <p>Freeze tube</p>	
 <p>Nasal swab</p>		 <p>Max 60 minutes</p>			 <p>Freeze tube upright</p>	
 <p>Sputum sample</p>		 <p>Max 60 minutes</p>			 <p>Freeze pot</p>	

Research Samples Visits 3, 4 & 5 (months 1, 3 & 6)

Tubes	Invert tube	Allow to stand	Centrifuge	Transfer	Store	Freeze at -80C
 <p>4-5 mls blood required.</p>	 <p>5 times</p>	 <p>Min 30 minutes Max 60 minutes</p>	 <p>1000g for 15 mins</p>	 <p>MAX 0.5ml per Fludix tube</p>	 <p>Up to 5 Fludix tubes</p>	
 <p>4 mls blood required</p>	 <p>8-10 times</p>	 <p>Max 60 minutes</p>			 <p>Freeze tube</p>	
 <p>Sputum sample</p>		 <p>Max 60 minutes</p>			 <p>Freeze pot</p>	